

XPOVIO[®]
(selinexor) tablets

Getting started with **XPOVIO**[®] (selinexor)

the first and only FDA-approved
oral medicine of its kind

What is XPOVIO?

XPOVIO[®] (selinexor) is a prescription medicine approved based on patient response rate. There are ongoing studies to confirm the clinical benefit of XPOVIO for the following use:

- to treat adults with certain types of diffuse large B-cell lymphoma that has come back (relapsed) or that did not respond to previous treatment (refractory) and who have received at least 2 prior therapies.

It is not known if XPOVIO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Your healthcare provider will do blood tests before you start taking XPOVIO and often during the first 3 months of treatment and then as needed during treatment.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).





IMPORTANT SAFETY INFORMATION, CONTINUED

XPOVIO can cause serious side effects, including:

- **Low platelet counts and low white blood cell counts are common with XPOVIO and can sometimes be severe or life-threatening.**
 - Low platelet counts can lead to bleeding, which can sometimes cause death. Your healthcare provider may prescribe platelet transfusions or other treatments for your low platelet counts.
Tell your healthcare provider right away if you have any bleeding or easy bruising during treatment with XPOVIO.
 - **Low white blood cell counts.** You may have an increased risk of infection during treatment with XPOVIO. Tell your healthcare provider if you have any signs or symptoms of infection, including fever, chills, cough, shortness of breath, pain during urination, or feeling generally unwell. If needed, your healthcare provider may prescribe antibiotics or certain medicines to help increase your white blood cell count.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO[®]
(selinexor) tablets

Table of contents

What is relapsed or refractory diffuse large B-cell lymphoma (RR DLBCL)?	4
Why XPOVIO® (selinexor)?	7
What is XPO1 and how does XPOVIO work to block it?	8
How did patients respond to treatment with XPOVIO?.....	10
What are the possible side effects with XPOVIO?	12
How do I take XPOVIO?	13
What else should I know about taking XPOVIO?	14
For caregivers: Caring for someone taking XPOVIO.....	16
What is KaryForward® and how do I join?	18
What additional support and resources are available?	19

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).



What is relapsed or refractory diffuse large B-cell lymphoma (RR DLBCL)?



Diffuse large B-cell lymphoma (DLBCL)

is a cancer that starts in B cells, a type of white blood cell. White blood cells are part of the immune system, which helps the body fight infection and disease. DLBCL can spread to other cells, organs, and tissues in the body.

RR DLBCL is cancer that has come back after treatment (relapse) or stopped responding to certain treatments (refractory).

RR DLBCL is treatable.

When DLBCL relapses, your doctor may start you on another type of treatment to help you get your disease under control.

How will my healthcare provider treat RR DLBCL?



There are different options for treating RR DLBCL, including:

Chemotherapy

Usually taken by mouth, injection, or infusion with or without other treatment to kill cancer cells

Stem cell transplants

Gives patients healthy stem cells (the blood-forming cells in the body) to replace their damaged cells

Cell therapies

A treatment in which T cells, a type of immune system cell, are collected from a patient's blood and modified in a laboratory. These modified cells can now help fight cancer cells and are returned to the patient's body via infusion.

Newer therapies

New types of medicines that fight RR DLBCL in different ways

How will my healthcare provider and I choose a treatment?



You and your healthcare provider will decide together which treatments are right for you

Some factors you may consider are:



How long the treatment has been shown to work in patients with RR DLBCL



Other health issues you may have



Possible **side effects** of treatment



Treatments you have taken before



If you are able to get to a clinic for your treatment

Why XPOVIO® (selinexor)?

XPOVIO is the first and only FDA-approved oral medicine of its kind that may treat RR DLBCL, even if other treatments no longer work for you


XPOVIO is a pill you take by mouth at home

You **DO NOT** have to go to the hospital to take XPOVIO

You **DO NOT** have to take XPOVIO with any other anti-cancer RR DLBCL medication
XPOVIO is targeted therapy, **NOT** chemotherapy

Kills at the cell core

XPOVIO works at the core of the cell (the nucleus) to kill some cancer cells



Responses were meaningful

29% of patients showed improvement with XPOVIO

The National Comprehensive Cancer Network® (NCCN) recommends selinexor (XPOVIO) for patients with RR DLBCL who have had at least 2 prior lines of systemic therapy and whose disease no longer responds to certain treatment options*

*Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for B-cell Lymphomas. V3.2025. © 2025 National Comprehensive Cancer Network, Inc. All rights reserved. Accessed October 28, 2025. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.


IMPORTANT SAFETY INFORMATION, CONTINUED

- **Serious infections.** XPOVIO can cause infections that can sometimes cause death, including upper or lower respiratory tract infections, such as pneumonia, and an infection throughout your body (sepsis). **Tell your healthcare provider right away if you have any signs or symptoms of an infection such as cough, chills, or fever during treatment with XPOVIO.**

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO®
(selinexor) tablets

What is XPO1 and why does it matter?




For illustrative purposes only.

XPO1 is a protein that maintains balance in healthy cells so they can function properly


- Cells have important anti-cancer materials that can only work in the cell's core (the nucleus)
- XPO1 is a protein that carries these materials from the nucleus to another part of the cell (the cytoplasm)
- To keep the cell in balance, XPO1 only carries the right amount of anti-cancer materials out of the nucleus

How does XPOVIO[®] (selinexor) work to block XPO1?

Some cancer cells have too much XPO1



In this cancer cell not exposed to XPOVIO




Cancer cell

XPO1 moves too many anti-cancer materials from the nucleus to the cytoplasm, so they can't work to fight the cancer.

Cancer cell growth

The cancer cells are able to grow and survive.

In this cancer cell exposed to XPOVIO



XPOVIO blocks XPO1 to kill the cancer cell at its core

In this cancer cell, XPOVIO stops XPO1 from carrying too many of the important anti-cancer materials out of the nucleus

- Anti-cancer materials stay in the nucleus
- Cancer cell dies

For illustrative purposes only.

IMPORTANT SAFETY INFORMATION, CONTINUED

- **Neurologic side effects.** XPOVIO can cause dizziness, fainting, decreased alertness, and changes in your mental status, including problems with thinking, seeing or hearing things that are not really there (hallucinations). These problems can sometimes be severe and life-threatening. Tell your healthcare provider right away if you get any of these symptoms and about all of the medicines you are taking. **Do not drive or operate heavy or dangerous machinery until you know how XPOVIO affects you. Take precautions to prevent a fall.**


Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO[®]
(selinexor) tablets



How did patients respond to treatment with XPOVIO® (selinexor)?

In the clinical trial, 134 adults with RR DLBCL were treated with XPOVIO 60 mg on Days 1 and 3 of every week. FDA approval was based upon the results of this trial, and additional trials may be needed.



29%
of patients
whose other treatments
stopped working showed
improvement when
treated with XPOVIO

Response to XPOVIO was evaluated in patients regardless of status (n=39):

- Their age
- The number of treatments they have taken before
- The type of DLBCL they have
- If DLBCL had stopped responding to other treatments
- If they had a stem cell transplant



XPOVIO has been shown to work when taken by itself in patients with RR DLBCL. Patients who responded did not need to take XPOVIO with another anti-cancer medicine.

IMPORTANT SAFETY INFORMATION, CONTINUED


- **Nausea, vomiting and/or diarrhea.** Nausea, vomiting and/or diarrhea can occur when you take XPOVIO and can sometimes be severe. You may be at risk for becoming dehydrated. Your healthcare provider may prescribe anti-nausea or anti-diarrhea medicines.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO[®]
(selinexor) tablets

How long did responses to XPOVIO® (selinexor) last?

Patients who responded to XPOVIO had a rapid and meaningful response



***The median duration of response (DOR) was a pre-specified secondary outcome, determined by a particular statistical analysis. Limitations to this analysis for the data from this study include a small number of patients and the effect of data from patients with prolonged remission durations. Approximately half of the responders in this analysis had a DOR lasting less than 3 months. The analysis of DOR is presented alongside an analysis demonstrating the number of patients responding at certain timepoints (3, 6, 12 months).**

The **median** number is the middle point of a set of numbers, in which half the numbers are above the median and half are below.

IMPORTANT SAFETY INFORMATION, CONTINUED

- **Loss of appetite and weight loss.** Loss of appetite and weight loss are common with XPOVIO. Tell your healthcare provider if you have a decrease or loss of appetite and if you are losing weight.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO®
(selinexor) tablets

What are the possible side effects with XPOVIO® (selinexor)?

Serious side effects

- Low platelet counts
- Low white blood cell counts
- Nausea, vomiting, and/or diarrhea
- Loss of appetite and weight loss
- Decreased sodium levels in your blood
- Serious infections
- Neurologic side effects
- New or worsening cataract, cloudiness or loss of transparency of the lens in the eye

Common side effects

- Tiredness
- Constipation
- Fever
- Low red blood cell count (anemia). Symptoms may include tiredness and shortness of breath.

These are not all the possible side effects of XPOVIO.

Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or [fda.gov/medwatch](https://www.fda.gov/medwatch).



In the clinical trial, the amount of time patients took XPOVIO ranged from 1 to 193 weeks, with the median amount of time being 9 weeks.

IMPORTANT SAFETY INFORMATION, CONTINUED

- **Decreased sodium levels in your blood.** Decreased sodium levels in your blood are common with XPOVIO but can sometimes be severe or life-threatening. Your healthcare provider may talk with you about your diet and prescribe IV fluids.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO®
(selinexor) tablets

How do I take XPOVIO® (selinexor)?

The recommended dosage of XPOVIO is 60 mg taken by mouth on Days 1 and 3 of each week



- Days 1 and 3 depend on which day of the week you start XPOVIO. For example, if you start XPOVIO on Tuesday, Tuesday is Day 1 and Thursday is Day 3
- You don't need to take XPOVIO with another anti-cancer medicine for RR DLBCL
- Take your medication as prescribed by your healthcare provider and be sure to let them know how you are feeling along the way
- Be sure to take any medicines prescribed by your healthcare provider before and during treatment with XPOVIO to help prevent nausea and vomiting. Tell your healthcare provider if the prescribed medicine does not control your nausea and vomiting

Tell your doctor if you have liver problems as that may impact your starting dose and/or dose adjustments.

IMPORTANT SAFETY INFORMATION, CONTINUED

- **New or worsening cataract, cloudiness or loss of transparency of the lens in the eye.** New or worsening cataract are common with XPOVIO. If a cataract forms, your vision may decrease, and you may need eye surgery to remove the cataract and restore your vision. **Tell your healthcare provider right away if you have symptoms of a cataract such as double vision, blurred vision, or sensitivity to light or glare.**

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).



During treatment, your healthcare provider may change your dose of XPOVIO. Always take XPOVIO exactly as prescribed.

XPOVIO[®]
(selinexor) tablets

What else should I know about taking XPOVIO® (selinexor)?



Talk to your healthcare provider before you change your dose or stop taking XPOVIO



Swallow XPOVIO whole with water. Never break, chew, crush, or divide the tablet



Take any medicines your healthcare provider prescribes for nausea/vomiting before and during treatment with XPOVIO



Tell your healthcare provider if the medicines don't help with nausea/vomiting



Drink enough fluids during treatment to prevent dehydration



Eat enough calories to prevent weight loss during treatment



If you miss a dose of XPOVIO, take your next dose at your next regularly scheduled day and time



If you vomit after taking a dose of XPOVIO, do not take an extra dose. Take your next dose at your next regularly scheduled day and time



If you take too much XPOVIO, call your healthcare provider right away



Do not share your XPOVIO prescription with other people



Your healthcare provider will do blood tests before you start taking XPOVIO, more often during the first 3 months of treatment, and then as needed

IMPORTANT SAFETY INFORMATION, CONTINUED

The most common side effects of XPOVIO in adults with DLBCL include:

- tiredness
- constipation
- fever
- low red blood cell count (anemia). Symptoms may include tiredness and shortness of breath.

Ability to have children: XPOVIO may affect the ability of both women and men to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all of the possible side effects of XPOVIO. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO[®]
(selinexor) tablets

What else should I ask my healthcare team before taking XPOVIO® (selinexor)?



- 1 Which **anti-nausea medicines** will I take prior to treatment with XPOVIO?
- 2 How often will I need a **blood test** while taking XPOVIO?
- 3 How do I ensure I **stay hydrated** while taking XPOVIO?
- 4 Should I expect to **lose weight** while taking XPOVIO? How much?
- 5 Will my **dose** be changed throughout treatment with XPOVIO?
- 6 How will my **side effects** be managed?
- 7 What should I do if I have a **new side effect or symptom**?
- 8 What should I know if I'm taking **other medications**?



Always ask your healthcare provider any questions you may have about treatment with XPOVIO.

IMPORTANT SAFETY INFORMATION, CONTINUED

Before taking XPOVIO, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had a recent or active infection
- have or have had bleeding problems
- have or have had liver problems
- have cataracts
- are pregnant or plan to become pregnant. XPOVIO can harm your unborn baby
- are taking prescription and over-the-counter medicines, vitamins, and herbal supplements

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

**XPOVIO**[®]
(selinexor) tablets



FOR CAREGIVERS:

How do I take care of myself when I'm caring for someone taking XPOVIO® (selinexor)?

Taking care of a loved one with RR DLBCL can be emotionally and physically demanding. It's important **not to lose sight of your own well-being**.

A support group that's online or in your local area may help you connect with people who understand what you're going through.

Your **loved one's healthcare providers and social worker** are there for you to answer your questions or support needs.

Doing activities like yoga and meditation may help relieve stress.

IMPORTANT SAFETY INFORMATION, CONTINUED

Females who are able to become pregnant: Your healthcare provider will check to see if you are pregnant before you start taking XPOVIO. You should use effective birth control (contraception) during treatment with XPOVIO and for 1 week after your last dose. Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with XPOVIO. Do not breastfeed during treatment with XPOVIO and for 1 week after your last dose of XPOVIO. It is not known if XPOVIO passes into your breast milk.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

XPOVIO[®]
(selinexor) tablets

FOR CAREGIVERS:

What else should I know about caring for someone taking XPOVIO® (selinexor)?



When caring for patients taking XPOVIO, keep in mind:



The **dosing schedule** for XPOVIO and anti-nausea medicines will be determined by the healthcare provider



Call the healthcare provider if your loved one experiences any side effects



The healthcare team will do **blood tests** before your loved one starts taking XPOVIO, more often during the first 3 months of treatment, and then as needed



Encourage your loved one to **drink lots of fluids and eat nutritious food** while on XPOVIO

IMPORTANT SAFETY INFORMATION, CONTINUED

Males with female partners who are able to become pregnant should use effective birth control (contraception) during treatment with XPOVIO and for 1 week after your last dose.

To report **SUSPECTED ADVERSE REACTIONS**, contact Karyopharm Therapeutics Inc. at 1-888-209-9326 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).

 **XPOVIO**[®]
(selinexor) tablets

What is KaryForward and how do I join?

KaryForward is a patient support program by Karyopharm Therapeutics dedicated to providing assistance and resources to you and your caregiver for XPOVIO® (selinexor)

KaryForward can help you with:



Insurance Coverage*

- Understand your benefits
- File claims
- Provide support during insurance appeals
- Find out what is covered by your insurance
- Access Karyopharm medications, even if there are unexpected delays or interruptions



Financial Assistance

- Qualify for the XPOVIO Copay Program and pay as little as \$5 for each XPOVIO prescription, or
- If eligible, get XPOVIO at no cost to you



Support and Resources

- Nurse Case Managers who can provide education and support during treatment
- Referrals to additional third-party support, such as transportation assistance

*Reimbursement Specialists are available to answer questions about the insurance approval process.

Please see Important Safety Information throughout, and [Medication Guide](#) and full [Prescribing Information](#).



**IF YOU'RE READY TO ENROLL
OR HAVE QUESTIONS ABOUT
KARYFORWARD:**

[CALL US](#)

1-877-KARY4WD (1-877-527-9493)

Monday through Friday, 8 AM to 8 PM ET

[VISIT US](#)

[KaryForward.com](https://www.KaryForward.com)



What additional support is available for my caregiver and me?

DLBCL Resources

Blood Cancer United

www.bloodcancerunited.org

800-955-4572

Lymphoma Research Foundation

www.lymphoma.org

800-500-9976

Lymphoma Coalition

www.lymphomacoalition.org

Caregiver Support Resources

Caregiver Action Network

www.caregiveraction.org

855-227-3640

Family Caregiver Alliance

www.caregiver.org

800-445-8106

General Cancer Resources

American Cancer Society

www.cancer.org

800-227-2345

CancerCare

www.cancercares.org

800-813-HOPE (4673)

Cancer Hope Network

www.cancerhopenetwork.org

877-HOPENET (467-3638)

Cancer Support Community

www.cancersupportcommunity.org

888-793-9355



In order to maintain the independence of these organizations, Karyopharm does not endorse or recommend any of the advocacy groups or services on this list and is not responsible for the content of these sites or services. Always talk to your healthcare team with any questions.

XPOVIO® (selinexor) is the first and only FDA-approved oral medicine of its kind that may treat RR DLBCL even if other treatments no longer work for you

- XPOVIO is a **pill you take by mouth** at home
- XPOVIO **works at the core of the cell** (the nucleus) to kill some cancer cells
- Patients who responded to treatment with XPOVIO were shown to have **a rapid and meaningful response**
- XPOVIO is targeted therapy, **not chemotherapy**

What is XPOVIO?

XPOVIO® (selinexor) is a prescription medicine approved based on patient response rate. There are ongoing studies to confirm the clinical benefit of XPOVIO for the following use:

- to treat adults with certain types of diffuse large B-cell lymphoma that has come back (relapsed) or that did not respond to previous treatment (refractory) and who have received at least 2 prior therapies.

It is not known if XPOVIO is safe and effective in children.

IMPORTANT SAFETY INFORMATION, CONTINUED

Your healthcare provider will do blood tests before you start taking XPOVIO and often during the first 3 months of treatment and then as needed during treatment.

Please see the [Medication Guide](#) and the full [Prescribing Information](#) for XPOVIO.

To report **SUSPECTED ADVERSE REACTIONS**, contact Karyopharm Therapeutics Inc. at 1-888-209-9326 or **FDA** at 1-800-FDA-1088 or www.fda.gov/medwatch.



All trademarks are the property of their respective owners.
© 2025 Karyopharm Therapeutics Inc. All rights reserved. US-XPOV-03/20-00041 (11/25)



Learn more at XPOVIO.com

**GET ASSISTANCE AND RESOURCES
FOR XPOVIO TREATMENT WITH
THE KARYFORWARD® PROGRAM.**

CALL

1-877-KARY4WD (1-877-527-9493)

Monday through Friday, 8 AM to 8 PM ET

VISIT

KaryForward.com

XPOVIO®
(selinexor) tablets